arXiv:1201.2041v2 [quant-ph] 14 Jan 2012

Monogamy of Measurement Induced Non-Locality

Ajoy Sen,[1,*] Debasis Sarkar,[1,†] and Amit Bhar[2,‡]
[1]*Department of Applied Mathematics, University of Calcutta, 92, A.P.C. Road, Kolkata-700009, India.*
[2]*Department of Mathematics, Jogesh Chandra Chaudhuri College,*
30, Prince Anwar Shah Road, Kolkata-700033, India.

(Dated: January 17, 2012)

Measurement Induced Non-Locality was introduced by Luo and Fu as a measure of non-locality in a bipartite state. In this paper we will discuss monogamy property of measurement induced non-locality for some three and four qubit classes of states. Unlike discord, we find quite surprising result in this situation. Both the generalized GHZ and W states satisfy monogamy relations in the three qubit case, however, in general there are violations of monogamy relations in both the GHZ-class and W-class states. In case of four qubit system, monogamy holds for most of the states in the generic class. Four qubit GHZ does not satisfy monogamy relation, but W-state does. We will also extend our results of generalized W-class to n-qubit.

PACS numbers: 03.67.Hk, 03.65.Ud, 03.65.Ta.

I. INTRODUCTION

Non-locality is in the heart of quantum world. From Bell's theorem[1], it is understood that no local hidden variable theory could replace quantum theory as a theory of physical world. The existence of entangled states in composite quantum systems assures the non-local behavior of quantum theory. Generally, violation of Bell's inequality is taken as the signature of non-locality. Entangled states play an important role to show the violation of Bell's inequality. However non-locality can be viewed from other perspectives. For example, there exist sets of locally indistinguishable orthogonal pure product states[2], used to show non-locality without entanglement. Recently introduced measurement induced non-locality[3] (in short, MIN), is one of the ways to detect non-locality in quantum states by some locally invariant measurements. Locally invariant measurements can not affect global states in classical theory but this is possible in quantum theory. So MIN is a type of non-local correlation which can only exists in quantum domain. MIN is defined in such a way that it is non-negative for all states, invariant under local unitary and vanishes on product state. In this sense, MIN could be observed as a type of non-local correlation which is induced by certain measurement. Although it is induced by some measurement, it is not a measure of quantum correlation in true sense. But it can be treated as a measure of non-locality induced by some kind of measurement.

Now a natural question arises about the shareability of quantum correlations in multipartite states. It may be monogamous or may be polygamous. It is known that entanglement is monogamous[4]. In this work we will investigate the monogamous behavior of MIN. We will check the monogamy property for some classes of states in both three and four-qubit system. Unlike discord[5–7], here we will show that both the three qubit generalized GHZ and W states satisfy monogamy relations, however there is violations of monogamy relation if we consider the generic whole class of pure three qubit states. In the case of pure four-qubit system we consider the most important generic class of states. It contains usual GHZ, maximally entangled states in the sense of Gour et.al. in[8]. Most of the states satisfy monogamy relations. There are two important subclasses of the generic class, say, $\mathcal{M}$ and τ_{min}. In one subclass monogamy holds but in another it does not. In particular, GHZ state violates monogamy relation and W satisfies it. Therefore, monogamous relations of MIN are quite different to that of some important measures of correlation[9, 10]. Our paper is organized as follows - in the section II we will review some of the basic properties of MIN. In section III we will explain and discuss about some four-qubit classes of states which we will require in our work. Section IV is devoted to the notion of monogamy for MIN. Section V contains results on pure three qubit systems and section VI contains results on four qubit system.

---

[*] Electronic address: ajoy.sn@gmail.com
[†] Electronic address: dsappmath@caluniv.ac.in
[‡] Electronic address: bhar.amit@yahoo.com

II. OVERVIEW ON MIN

Consider ρ be any bipartite state shared between two parties A and B. Then MIN (denoted by $N(\rho)$) is defined by,

$$N(\rho) := \max_{\{\Pi^A\}} \parallel \rho - \Pi^A(\rho) \parallel^2 \tag{1}$$

where maximum is over all Von Neumann measurements $\{\Pi^A\}$ which do not disturb ρ_A, the local density matrix of A, i.e., $\Sigma_k \Pi_k^A \rho_A \Pi_k^A = \rho_A$ and $\|.\|$ is taken as Hilbert Schmidt norm (i.e. $\parallel X \parallel = [Tr(X^\dagger X)]^{\frac{1}{2}}$). It is in some sense, dual to that of geometric measure of discord. Physically, MIN quantifies the global effect caused by locally invariant measurements. MIN has application in general dense coding, quantum state steering etc. MIN vanishes for product state and remains positive for entangled states. For pure states MIN reduces to linear entropy like geometric discord[9]. Explicit formula of MIN for $2 \otimes n$ system, $m \otimes n$ (if ρ_A is non-degenerate) system and an explicit upper bound for $m \otimes n$ system were obtained by Luo, Fu in [3]. Later Mirafzali *et.al.* [11] formulate a way to reduce the problem of degeneracy in $m \otimes n$ system and evaluate it for $3 \otimes n$ dimensional system. MIN is invariant under local unitary, i.e., in true sense, it is a non-local correlation measure. The set of all zero MIN states is non-convex. Guo, Hou [12] derived the conditions for the nullity of MIN. They have found that set of states with zero MIN is a proper subset of the set of all classical quantum states, i.e., zero discord states. MIN for classical quantum state vanishes if each eigen-subspace of ρ_A is one dimensional. It therefore reveals that non-commutativity is the cause of these kind of non-locality in quantum states. Recently, in [13], MIN have been quantified in terms of relative entropy to give it another physical interpretation. However, in our work we have used the older definition of MIN.

Suppose H^A, H^B be the Hilbert spaces associated with the parties A and B respectively and $L(H^A)$, $L(H^B)$ denote the Hilbert space of linear operators acting on H^A, H^B with the inner product defined by $\langle X|Y\rangle := trX^\dagger Y$. We state two important results which we will use in our work.

Theorem 1:(Luo, Fu [3]) *Let* $|\psi\rangle_{AB}$ *be any bipartite pure state with Schmidt decomposition* $|\psi\rangle_{AB} = \sum_i \sqrt{s_i}|\alpha_i\rangle_A|\beta_i\rangle_B$, *then* $N(|\psi\rangle_{AB}) = 1 - \sum_i s_i^2$.

Theorem 2:(Luo, Fu [3]) *If* ρ_{AB} *be any state of* $2 \otimes n$ *dimensional system written in the form*

$$\rho_{AB} = \frac{1}{\sqrt{2n}} \frac{I^A}{\sqrt{2}} \otimes \frac{I^B}{\sqrt{n}} + \sum_{i=1}^{3} x_i X_i^A \otimes \frac{I^B}{\sqrt{n}} + \frac{I^A}{\sqrt{2}} \otimes \sum_{i=1}^{n^2-1} y_i Y_i^B + \sum_{i=1}^{3}\sum_{j=1}^{n^2-1} t_{ij} X_i^A Y_j^B$$

where $\{X_i^A : i=0,1,2,3\}$ *and* $\{Y_j^B : j = 0, 1, 2..n^2 - 1\}$ *are the orthonormal hermitian operator bases for* $L(H^A)$ *and* $L(H^B)$ *respectively with* $X_0^A = I^A/\sqrt{2}, Y_0^B = I^B/\sqrt{n}$. *Then*

$$N(\rho_{AB}) = \begin{cases} trTT^t - \frac{1}{\|\boldsymbol{x}\|^2}\boldsymbol{x}^t TT^t \boldsymbol{x} & \text{if } \boldsymbol{x} \neq \boldsymbol{0} \\ trTT^t - \lambda_3 & \text{if } \boldsymbol{x} = \boldsymbol{0} \end{cases} \tag{2}$$

where the matrix $T = (t_{ij})_{3\times(n^2-1)}$ *with* λ_3 *being minimum eigenvalue of* TT^t *and* $\|\boldsymbol{x}\|^2 := \sum_i x_i^2$ *with* $\boldsymbol{x} = (x_1, x_2, x_3)^t$.

Before going to discuss monogamy properties of measurement induced non locality, specifically for three and four qubit systems, we first mention some important classes of states in four qubit systems with some discussions on their entanglement behavior.

III. SOME SPECIAL FOUR QUBITS CLASSES

Four qubit pure states can be classified in nine groups[14]. Among them the generic class is given by

$$\mathcal{A} \equiv \{\sum_{j=0}^{3} z_j u_j : \sum_{j=0}^{3} |z_j|^2 = 1, z_i \in \mathbb{C}, i = 0, 1, 2, 3\} \tag{3}$$

where $u_0 \equiv |\phi^+\rangle|\phi^+\rangle$, $u_1 \equiv |\phi^-\rangle|\phi^-\rangle$, $u_2 \equiv |\psi^+\rangle|\psi^+\rangle$ and $|\phi^\pm\rangle = (|00\rangle \pm |11\rangle)/\sqrt{2}$, $|\psi^\pm\rangle = (|01\rangle \pm |10\rangle)/\sqrt{2}$. Consider two important subclasses $\mathcal{M}$ and τ_{min} of the generic class $\mathcal{A}$ which are defined as,

$$\mathcal{M} = \{\sum_{j=0}^{3} z_j u_j : \sum_{j=0}^{3} |z_j|^2 = 1, \sum_{j=0}^{3} z_j^2 = 0\} \tag{4}$$

and

$$\tau_{min} = \{\sum_{j=0}^{3} x_j u_j : \sum_{j=0}^{3} x_j^2 = 1, x_j \in \mathbb{R}, j = 0,1,2,3\} \tag{5}$$

These two subclasses are important in the sense that $\mathcal{M}$ is the maximally entangled class and τ_{min} have least amount of bipartite entanglement according to the definition of maximally entangled states, given by Gour et. al. in [8]. Consider a pure bipartite state $|\psi_{AB}\rangle \in \mathbb{C}^m \otimes \mathbb{C}^n$. Then the tangle is defined as

$$\tau_{AB} = 2(1 - tr\rho_A^2) \tag{6}$$

where $\rho_A = Tr_B|\psi_{AB}\rangle\langle\psi_{AB}|$. Now for a pure state shared between four parties $|\psi_{ABCD}\rangle$, three quantities $\tau_1, \tau_2, \tau_{ABCD}$ are defined as:

$$\tau_1 \equiv \frac{1}{4}(\tau_{A|BCD} + \tau_{B|ACD} + \tau_{C|ABD} + \tau_{D|ABC}) \tag{7}$$

$$\tau_2 \equiv \frac{1}{3}(\tau_{AB|CD} + \tau_{CA|BD} + \tau_{DA|BC}) \tag{8}$$

$$\tau_{ABCD} = 4\tau_1 - 3\tau_2 \tag{9}$$

For the above two subclasses $\mathcal{M}$ and τ_{min} we have $\tau_1(\mathcal{M}) = 1$, $\tau_2(\mathcal{M}) = \frac{4}{3}$, $\tau_{ABCD}(\mathcal{M}) = 0$ and $\tau_1(\tau_{min}) = 1$, $\tau_2(\tau_{min}) = 1$, $\tau_{ABCD}(\tau_{min}) = 1$. Four-qubit GHZ state belongs to the class τ_{min}.

IV. MONOGAMY

Monogamy is an important aspect in our physical world which restricts on the shareability of bipartite correlation. Entanglement is an example of quantum correlation which is monogamous w.r.t. the tangle[4]. Mathematically a correlation measure Q is said to be monogamous iff for any n-party state $\rho_{A_1A_2..A_n}$ the relation

$$\sum_{k=1,k\neq i}^{n} Q(\rho_{A_iA_k}) \leq Q(\rho_{A_i|A_1A_2..A_{i-1}A_{i+1}..A_n}) \tag{10}$$

holds for all $i = 1,2...n$. Now consider an n-party state $\rho_{12...n}$. Let the locally invariant measurement is done on the party 1. Then MIN is defined as $N(\rho_{1|2...n}) =\| \rho_{12...n} - \Pi_1^*(\rho_{1|2...n}) \|^2$, where $\Pi_1^* = \{\pi_{1k}^*\}$ is the optimal measurement done by the party 1 which does not change its local density matrix, i.e., $\rho_1 = \Sigma_k \pi_{1k}^* \rho_1 \pi_{1k}^*$. On the other hand, since $\rho_1 = Tr_{23..n}(\rho_{12..n}) = Tr_j(Tr\rho_{1j}), j = 2,3..n$, the optimal measurement also does not change the local density matrices for all two-party reduced states of $\rho_{12...n}$ of the kind, $\rho_{1j} = Tr_{2,3..j-1,j+1...n}(\rho_{1,2...n})$. Then $\Sigma_j N(\rho_{1j}) \geq \sum_j \|\rho_{1j} - \Pi_1^*(\rho_{1j})\|^2$. So in case of polygamy $N(\rho_{1|2...n}) < \sum_j \|\rho_{1j} - \Pi_1^*(\rho_{1j})\|^2$ and if $N(\rho_{1|2...n}) \geq \sum_j \|\rho_{1j} - \Pi_1^*(\rho_{1j})\|^2$ then the state is monogamous w.r.t. MIN.

V. MONOGAMY IN THREE QUBIT SYSTEM

Any three qubit pure state $|\psi_{ABC}\rangle$ has a generic form $\lambda_0|000\rangle + \lambda_1 e^{i\theta}|100\rangle + \lambda_2|101\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle$, where $\lambda_i \in \mathbb{R}, \theta \in [0,\pi], \sum_i \lambda_i^2 = 1$[15] . This class includes GHZ class. W class can also be availed by putting $\lambda_4 = 0$. For the general state $|\psi_{ABC}\rangle$, the reduced density matrix $\rho_{AB} = Tr_C|\psi_{ABC}\rangle\langle\psi_{ABC}|$ has the form

$$\rho_{AB} = \begin{bmatrix} \lambda_0^2 & 0 & \lambda_0\lambda_1 e^{-i\theta} & \lambda_0\lambda_3 \\ 0 & 0 & 0 & 0 \\ \lambda_0\lambda_1 e^{i\theta} & 0 & \lambda_1^2 + \lambda_2^2 & \lambda_1\lambda_3 e^{i\theta} + \lambda_2\lambda_4 \\ \lambda_0\lambda_3 & 0 & \lambda_1\lambda_3 e^{-i\theta} + \lambda_2\lambda_4 & \lambda_3^2 + \lambda_4^2 \end{bmatrix}$$

Then the correlation matrix $T = (t_{ij})_{3\times3}$, as obtained from the relation $t_{ij} = tr(\rho\frac{\sigma_i}{\sqrt{2}} \otimes \frac{\sigma_j}{\sqrt{2}}), i,j = 1,2,3; \sigma_i$'s being the Pauli matrices, is

$$T = \begin{bmatrix} \lambda_0\lambda_3 & 0 & \lambda_0\lambda_1\cos\theta \\ 0 & -\lambda_0\lambda_3 & -\lambda_0\lambda_1\sin\theta \\ -\lambda_1\lambda_3\cos\theta - \lambda_2\lambda_4 & -\lambda_1\lambda_3\sin\theta & 0.5 - \lambda_1^2 - \lambda_3^2 \end{bmatrix}$$

Other reduced density matrix ρ_{AC} and its corresponding correlation matrix could be written at once from ρ_{AB} and T by only inter changing λ_2 and λ_3. Coherent vector for both the reduced density matrices is $\mathbf{x} = (\lambda_0\lambda_1\cos\theta, -\lambda_0\lambda_1\sin\theta, \lambda_0^2 - 0.5)^t$.
Clearly, $\parallel \mathbf{x} \parallel = 0$ iff $\lambda_0^2 = 0.5, \lambda_1^2 = 0$. In case of $\parallel \mathbf{x} \parallel = 0$ we have,

$$N(\rho_{AB}) = 2a + c - \min\{a, \frac{1}{2}(a + c - \sqrt{(a-c)^2 + 4b^2})\} \tag{11}$$

$$N(\rho_{AC}) = 2g + k - \min\{g, \frac{1}{2}(g + k - \sqrt{(g-k)^2 + 4b^2})\} \tag{12}$$

$$N(\rho_{A|BC}) = 0.5 \tag{13}$$

where $a = \lambda_0^2\lambda_3^2, b = -\lambda_0\lambda_2\lambda_3\lambda_4, c = \lambda_2^2\lambda_4^2 + (0.5 - \lambda_3^2)^2, g = \lambda_0^2\lambda_2^2, k = \lambda_3^2\lambda_4^2 + (0.5 - \lambda_2^2)^2$ with $\lambda_0^2 = 0.5$. Now the minimum value of $N(\rho_{AB}) + N(\rho_{AC})$ is 0.5. So in case of $\parallel \mathbf{x} \parallel = 0$ monogamy is violated for most of the states. Numerical simulation of 10^6 states shows that around 0.02% of three qubit generic states and around 20% W class states satisfy monogamy.
When $\parallel \mathbf{x} \parallel \neq 0$ we have

$$N(\rho_{A|BC}) = 2\lambda_0^2(\lambda_2^2 + \lambda_3^2 + \lambda_4^2) \tag{14}$$

$$N(\rho_{AB}) = a + b + c - \frac{1}{\|\mathbf{x}\|^2}\mathbf{x}^t T T^t \mathbf{x} \tag{15}$$

$$N(\rho_{AC}) = g + f + k - \frac{1}{\|\mathbf{x}\|^2}\mathbf{x}^t T T^t \mathbf{x} \tag{16}$$

where $a = \lambda_0^2\lambda_3^2 + \lambda_0^2\lambda_1^2\cos^2\theta$, $b = \lambda_0^2\lambda_3^2 + \lambda_0^2\lambda_1^2\sin^2\theta$, $c = (\lambda_2\lambda_4 + \lambda_1\lambda_3\cos\theta)^2 + \lambda_1^2\lambda_3^2\sin^2\theta + (0.5 - \lambda_1^2 - \lambda_3^2)^2$, $g = \lambda_0^2\lambda_2^2 + \lambda_0^2\lambda_1^2\cos^2\theta$, $f = \lambda_0^2\lambda_2^2 + \lambda_0^2\lambda_1^2\sin^2\theta$, $k = (\lambda_3\lambda_4 + \lambda_1\lambda_2\cos\theta)^2 + \lambda_1^2\lambda_2^2\sin^2\theta + (0.5 - \lambda_1^2 - \lambda_2^2)^2$. Now the maximum value of $N(\rho_{AB}) + N(\rho_{AC})$ is 0.5. Hence three qubit pure states with $\parallel \mathbf{x} \parallel \neq 0$ satisfies the monogamy relation. Specifically, the three qubit generalized GHZ class of pure states($\alpha|000\rangle + \beta|111\rangle$) is monogamous in the region $\alpha \neq \beta$ (equivalent to $\parallel \mathbf{x} \parallel \neq 0$)) and the monogamy relation holds good with equality if $\alpha = \beta (= \frac{1}{\sqrt{2}})$ (equivalent to $\parallel \mathbf{x} \parallel = 0$). In the three qubit generalized W-Class states($\alpha|001\rangle + \beta|010\rangle + \gamma|001\rangle$) we have $N(\rho_{AB}) + N(\rho_{AC}) = N(\rho_{A|BC}) = 2|\alpha|^2(1 - |\alpha|^2)$. That is the monogamy relation holds with equality.

VI. MONOGAMY IN FOUR QUBIT SYSTEM

Consider a four-qubit generic pure state $|\psi_{ABCD}\rangle$ which is shared between four parties A, B, C, D (i.e.,$|\psi_{ABCD}\rangle \in \mathcal{A}$) where

$$|\psi_{ABCD}\rangle = \sum_{j=0}^{3} z_j u_j, \quad \sum_{j=0}^{3} |z_j|^2 = 1. \tag{17}$$

We consider the two-qubit reduced density matrices ρ_{AB}, ρ_{AC} and ρ_{AD} of $\rho = |\psi_{ABCD}\rangle\langle\psi_{ABCD}|$. Each reduced density matrix is of the form

$$\frac{1}{4}\begin{bmatrix} \alpha & 0 & 0 & \beta \\ 0 & \gamma & \delta & 0 \\ 0 & \delta & \gamma & 0 \\ \beta & 0 & 0 & \alpha \end{bmatrix}$$

where $\alpha, \beta, \gamma, \delta$ are some suitable functions of z_0, z_1, z_2, z_3 such that $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and $\alpha + \gamma = 2$. We define 4 quantities $a = z_0 + z_1, b = z_0 - z_1, c = z_2 + z_3, d = z_2 - z_3$. Then for ρ_{AB}: $\alpha = 2(|z_0|^2 + |z_1|^2), \beta = 2(|z_0|^2 - |z_1|^2), \gamma = 2(|z_2|^2 + |z_3|^2), \delta = 2(|z_2|^2 - |z_3|^2)$; for ρ_{AC}: $\alpha = (|a|^2 + |c|^2), \beta = 2Re(\overline{a}c), \gamma = 2(|b|^2 + |d|^2), \delta = 2Re(\overline{b}d)$ and for ρ_{AD}: $\alpha = (|a|^2 + |d|^2), \beta = 2Re(\overline{a}d), \gamma = 2(|b|^2 + |c|^2), \delta = 2Re(\overline{b}c)$. The elements of the correlation matrix T can be obtained from $t_{ij} = tr(\rho_{AX} X_i \otimes Y_j),\ X = B, C, D$. Eigenvalues of the matrix TT^t are of the form $k(\beta \pm \delta)^2, k(\alpha - \gamma)^2$ with $k = \frac{1}{16}$. The coherent vector $\mathbf{x} = (x_1, x_2, x_3)^t$ as obtained from the relation $x_i = tr(\rho_{AX} X_i \otimes I),\ X = B, C, D$ is zero for all the three cases. Hence we have(by theorem 2),

$$N(\rho_{AX}) = k.\ [2(\beta^2 + \delta^2) + (\alpha - \gamma)^2] - \lambda_3 \quad where \quad X = B, C, D \tag{18}$$

and

$$\lambda_3 = k.\ min\{(\beta + \delta)^2, (\alpha - \gamma)^2, (\beta - \delta)^2\}$$

On the other hand, we can write the state $\rho_{A|BCD}$ in the form $\frac{1}{\sqrt{2}}(|0\phi_0\rangle + |1\phi_1\rangle)$ where $|\phi_0\rangle, |\phi_1\rangle$ are mutually orthonormal. Since this is a pure state we have $N(\rho_{A|BCD}) = 0.5$ (using theorem 1). Numerical simulation for 10^6 generic states shows that about 66% of them satisfies monogamy relation. Hence in general four-qubit generic class is not monogamous w.r.t. MIN. So there exist quantum state whose locally shared non-locality exceeds the amount of globally shared non-locality.



10^5 *simulation shows about 34% violation of monogamy for the generic class* $\mathcal{A}$*. The red line marks the departure from monogamy.*

However for the sub classes $\mathcal{M}$ and τ_{min} we can still check whether monogamy relation holds or not and if not then what is the amount of violation. These things will be clear in the next few results.

Theorem 3: *Consider a four-qubit system.Then for any pure state* ρ *belonging to the generic class* $\mathcal{A}$ *we have* $N(\rho_{XY}) \leq (\sum_{i=1}^{3} \lambda_i)$ *where* ρ_{XY} *denotes any bipartite reduced density matrix of* ρ *and* λ_i*'s are the eigenvalues of* TT^t*,* T *being the correlation matrix of* ρ_{XY}*.*

Proof: The proof of the theorem is very easy as it directly follows form Theorem 2 keeping in mind that the eigenvalues of TT^t are all positive and the Bloch vector $\|\mathbf{x}\| = \mathbf{0}$ for all bipartite reduced states of any generic state. We will use this theorem in our proof of the the next two results. □

Theorem 4: *Let* ρ_{ABCD} *be any four-qubit pure state belonging to the class* $\mathcal{M}$*. Then* $N(\rho_{AB}) + N(\rho_{AC}) + N(\rho_{AD}) \leq \frac{1}{4}$

Proof: Let us consider any four qubit pure state $\rho_{ABCD} = |\psi\rangle_{ABCD}\langle\psi|$ belonging to the class $\mathcal{M}$ i.e., $|\psi\rangle_{ABCD} = \sum_{j=0}^{3} z_j u_j$, with $\sum_{j=0}^{3} |z_j|^2 = 1, \sum_{j=0}^{3} z_j^2 = 0$. Then, taking $z_j = x_j + i.y_j\ \forall j = 0, 1, 2, 3; x_j, y_j \in \mathbb{R}$ and utilizing the above restrictions, we get $\sum_{j=0}^{3} x_j^2 = \sum_{j=0}^{3} y_j^2 = \frac{1}{2}$ and $\sum_{j=0}^{3} x_j y_j = 0$. The fruitful implementation of the result of previous theorem and simple algebraic manipulations using these results lead to the fact $N(\rho_{AB}) + N(\rho_{AC}) + N(\rho_{AD}) \leq \frac{1}{4}$.□

Theorem 5: *Let* ρ_{ABCD} *be any four-qubit pure state belonging to the class* τ_{min}*.Then* $\frac{1}{2} \leq N(\rho_{AB}) + N(\rho_{AC}) + N(\rho_{AD}) \leq \frac{3}{4}$.

Proof: Let us consider any four-qubit pure state $\rho_{ABCD} = |\psi\rangle_{ABCD}\langle\psi|$, belonging to the class τ_{min} i.e., $|\psi\rangle_{ABCD} = \sum_{j=0}^{3} x_j u_j$, with $\sum_{j=0}^{3} x_j^2 = 1$ and $x_j \in \mathbb{R}, j = 0, 1, 2, 3$. Now let us denote the sets of eigenvalues of ρ_{AB}, ρ_{AC} and ρ_{AD} as $\{\lambda_1^{AB}, \lambda_2^{AB}, \lambda_3^{AB}\}$, $\{\lambda_1^{AC}, \lambda_2^{AC}, \lambda_3^{AC}\}$ and $\{\lambda_1^{AD}, \lambda_2^{AD}, \lambda_3^{AD}\}$ respectively. Without loss of generality we can consider, $\lambda_i^{AB} \geq \lambda_j^{AB} \geq \lambda_k^{AB}, \forall i, j, k = 1, 2, 3$ with $i \neq j \neq k$. This above consideration gives natural connections among the eigenvalues of ρ_{AC} and also ρ_{AD}. These are $\lambda_i^{AC} \geq \lambda_j^{AC} \geq \lambda_k^{AC}$ and $\lambda_i^{AD} \geq \lambda_j^{AD} \geq \lambda_k^{AD}$, $\forall i, j, k = 1, 2, 3$ with $i \neq j \neq k$. Observing the behavior of the eigenvalues and straight forward derivation quite easily establish the desired result, i.e., $\frac{1}{2} \leq N(\rho_{AB}) + N(\rho_{AC}) + N(\rho_{AD}) \leq \frac{3}{4}$.□

These two results give us information about the monogamous behavior of MIN in two important classes of four-qubit system. Since, $N(\rho_{A|BCD}) = \frac{1}{2}$ in the whole class $\mathcal{A}$, therefore, MIN is monogamous in the subclass $\mathcal{M}$ but it is polygamous in other class τ_{min}. Further, all the states that are connected to these classes by LU, share the same fate. Four-qubit GHZ state belongs to the class τ_{min}. Hence GHZ and their LU equivalent states are not monogamous w.r.t MIN. Another two states(and obviously their LU equivalent states) $|L\rangle = \frac{1}{\sqrt{3}}(u_0 + \omega u_1 + \omega^2 u_2)$ where $\omega = e^{\frac{2i\pi}{3}}$ and $|M\rangle = \frac{i}{\sqrt{2}}u_0 + \frac{1}{\sqrt{6}}(u_1 + u_2 + u_3)$ which maximize the Tsallis α-entropy for different regions of α [8], satisfy the monogamy relation of MIN. On the other hand the four-qubit cluster states satisfies the monogamy relation as their two party reduced density matrices are completely mixed (i.e., MIN is zero). For four-qubit generalized W-states $\rho^W = \alpha|1000\rangle + \beta|0100\rangle + \gamma|0010\rangle + \delta|0001\rangle$ where $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$, monogamy relation holds with equality, i.e., it can be easily shown (as in the three-qubit case) that $N(\rho^W_{AB}) + N(\rho^W_{AC}) + N(\rho^W_{AD}) = N(\rho^W_{A|BCD}) = 2|\alpha|^2(1 - |\alpha|^2)$. Hence for this type of states non-locality shows additive property with respect to each party. This result of generalized W- class can be further extended to n-qubit system with the same conclusion.

VII. CONCLUSION

Thus we have explored the monogamy nature of MIN and found certain classes of states on which MIN shows monogamy relations. Unlike geometric discord,[7] MIN can be polygamous for pure state as revealed in some subclass of generic class. On the other hand, W-class seems to satisfy the monogamy relation with equality in n-qubit system. So for the W-class MIN becomes additive in terms of sharing between the parties. Thus existence of Monogamy of this type of correlation put a restriction on the amount of shared non-locality.

Acknowledgement. The author A. Sen acknowledges the financial support from University Grants Commission, New Delhi, India.

[1] J.S. Bell, Physics 1, 195,(1964); Speakable and Unspeakable in Quantum Mechanics, Cambridge University Press (1987).

[2] Charles H. Bennett, David P. DiVincenzo, Christopher A. Fuchs, Tal Mor, Eric Rains, Peter W. Shor, John A. Smolin, and William K. Wootters, Phys. Rev. A 59, 1070 (1999).

[3] S. Luo and S. S. Fu, Phys. Rev. Lett. 106, 120401 (2011).

[4] Valerie Coffman, Joydip Kundu, William K. Wootters, Phys. Rev. A 61,052306 (2000).

[5] Gian Luca Giorgi, Phys. Rev. A 84, 054301 (2011).

[6] R. Prabhu, Arun Kumar Pati, Aditi Sen(De), and Ujjwal Sen, arXiv: 1108.5168v1.

[7] Alexander Streltsov, Gerardo Adesso, Marco Piani and Dagmar Bruss, arXiv: 1112.3967v1.

[8] G. Gour and N. R. Wallach, J. Math. Phys. 51(11), 112201 (2010).

[9] Borivoje Dakic,Vlatko Vedral and Caslav Brukner,Phys. Rev. Lett. 105, 190502 (2010).

[10] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. 88, 017901 (2001).

[11] S. Y. Mirafzali, I. Sargolzahi, A. Ahanj, K. Javidan, and M. Sarbishaei, arXiv:1110.3499.

[12] Yu Guo, and Jinchuan Hou, arXiv:1107.0355v1.

[13] Zhengjun Xi, Xiaoguang Wang and Yongming Li, arXiv:1112.0736v1.

[14] F. Verstraete, J. Dehaene, B. De Moor, H. Verschelde, Phys. Rev. A 65, 052112 (2002).

[15] A. Acin, A. Andrianov, E. Jane, R. Tarrach, J. Phys. A: Math. Gen. 34 (2001), 6725.